

CAL 40

NPKS 7-0-0-0 + 40% Calcium



High analysis suspension for supplying calcium, correction of calcium deficiencies & improving shelf life and storage quality of produce.

BENEFITS OF CAL 40

- Highly micronised, controlled released, low salt index calcium. Suitable for all crops.
- With a high analysis it contains more than twice the calcium found in Cal Nitrate.
- Controlled particle morphology to delivering sustained uptake over 4 – 6 weeks.
- Provides sunburn mitigation via the formation of a particulate barrier.
- Enhances soil health and rejuvenation by improving pH.
- Improves the soil structure by displacing sodium and chloride thereby improving the soil water infiltration and aeration.

THE IMPORTANCE OF CALCIUM AND NITROGEN

Calcium is required for the cellulose precursors in cell wall formation. It also stabilises cell membranes and protects them, an important attribute under stress conditions. In fruit crops it is required in high quantities as it is important for fruit quality and shelf life. When plants are threatened by infection, calcium combines with a protein to stimulate salicylic acid (SA) production. Calcium deficiency leads to poor fruit set, blossom-end-rot, bitter pit, cell collapse and tissue death.

THE IMPORTANCE OF NITROGEN

CAL 40 helps improve the soil PH from acidic range to neutral upon consistent use. If blended with ACTIVIST® MAG-FLO the micro-fine particles of CAL 40 and ACTIVIST® MAG-FLO neutralise the aluminum and iron phosphates in acid soils.

CAL 40

CHARACTERISTICS: pH: 8.5 – 10; Specific Gravity: 1.70 – 1.73

AUS Analysis W/W%: 6.9% N, 40% Ca.

International Analysis W/W%: 4.0% N, 23.3% Ca.

APPLICATION

BROADACRE: Such as Barley, Canola, Cotton, Grain legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture crops. **Foliar: 1.5 – 3 L/ha** in a minimum of 30 – 60L final spray volume. Foliar Spray, early tillering to jointing stage.

DECIDUOUS TREE CROPS: Such as Apple, Almond, Cherry, Nectarine, Peach, Pear, Pistachio and Walnut. **Foliar: 4 – 7 L/ha** in a minimum of 600 – 1000L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Apply at early spur burst, complete petal fall and post blossom as required. **DO NOT apply as foliar on high chill stone fruit varieties.**

EVERGREEN TREE CROPS: Such as Avocado, Citrus, Macadamia, Lychee. **Foliar: 4 – 7 L/ha** in a minimum of 400 – 700L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Apply at flower bud break and spring flush with follow-up applications through fruit fill as required. Note: Do Not apply later than 6 weeks prior to harvest as residue may remain.

FRUITING VEGETABLES: Such as Capsicum, Cucurbits, Eggplant, Tomatoes (field), Watermelons, Pumpkins. **Foliar: 3 – 6 L/ha** in a minimum of 300 – 600L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Apply when plants are commencing flowering and repeat at 10 - 14 day intervals, or as required.

LEAFY VEGETABLES: Such as Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale and Herbs. **Foliar: 3 – 5 L/ha** in a minimum of 300 – 500L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Foliar spray 10 – 14 days post transplant.

ROOT VEGETABLES: Such as Beetroot, Carrot, Leek, Onion, Potato, Radish, Sweet Potato. **Foliar: 3 - 5 L/ha** in a minimum of 300 - 500L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Apply when sufficient leaf area present, repeat every 3 - 4 weeks. Potatoes: After emergence and during canopy closure, fertigate at bulking.

VINE and BERRY CROPS: Such as Blueberry, Strawberry, Raspberry, Wine and Table Grapes. **Foliar: 2.5 – 4 L/ha** in a minimum of 250 – 400L final spray volume. **Fertigation: 10 – 40 L/ha**, Higher rates are for pH adjustment. Foliar spray 3 treatments, shoots 10cm, flower buds separated & fruit set. For table grapes last treatment to be 1 month prior to harvest. Use double rate post harvest before leaf fall to improve nutrient levels prior to dormancy.

Fertigation rates are dependent on seasonal nutrient demand.

Agitate contents well prior to application.

DO NOT APPLY to high pH soil via fertigation. Foliar applications are preferred.

The information contained in this Product Information Sheet in respect of the "Product" is indicative only and should not be relied upon as advice or a recommendation. While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information. You use the information at your own risk and should rely on your own independent inquiries and assessments. With the exception of the consumer guarantees provided by the Australian Consumer Law (ACL), all conditions and warranties implied in respect of any information or advice provided by Agrichem about the Product are excluded, and Agrichem does not accept any liability whatsoever (including through misrepresentation or negligence), incurred in connection with your use or reliance upon this Information Sheet. If liability under the ACL cannot be excluded but the Product the subject of the Information Sheet is NOT used for personal, domestic or household use or consumption, Agrichem may (at its election) limit its liability to replacement of the Product, or payment of the cost of acquiring the Product. You must not reproduce this information sheet without written consent from Agrichem®.

NOTE: The suggested rates of application of the Product are designed for typical Australian conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.