

AGRI K 415

NPKS 0 - 0 - 41.5 - 0



Enhanced availability potassium for improving colour, size, ripening and flavour of fruit and vegetables

BENEFITS OF AGRI K 415

- Highly concentrated, enhanced availability potassium (K).
- Provides more K for your crop quickly to maximize plant strength & disease resistance.
- Formulated for maximum plant uptake & crop safety.
- Convenient liquid formulation.
- Improves in-plant sugar production and enhances fruit colour and size.

THE IMPORTANCE OF POTASSIUM

Potassium optimises water use efficiency and is the key nutrient to improve crop photosynthesis and sugar production in fruits. Potassium is very important in fruit bearing plants. Potassium regulates the electrolytes and turgidity of plant cells. Potassium occurs in the guard cells of the stomata and is therefore essential in respiration and transpiration. Potassium is required at all growth stages and a lack of potassium cannot be rectified with late applications.

AGRI K 415

CHARACTERISTICS: pH: 13; Specific Gravity: 1.515 – 1.525

AUS Analysis W/V%: 41.5% K,
International Analysis W/W%: 32.9% (K₂O)

APPLICATION

BROADACRE: Such as Barley, Canola, Cotton, Grain legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture crops. **Foliar: 3 – 6 L/ha** in a minimum of 60 - 120 L final spray volume. Canola: Foliar spray at growth stage one - 4 or more leaves. Repeat at onset of stem elongation. Cereals: Foliar spray four – five leaf to early stem extension Zodok's G.S. 12 - 30.

DECIDUOUS TREE CROPS: Such as Apple, Almond, Cherry, Nectarine, Peach, Pear, Pistachio and Walnut. **Fertigation: 10 – 20 L/ha.** Apply as required to encourage & maintain nutrient levels. Note: **DO NOT** apply as a foliar to stone fruit during leaf growth. Can be applied Post harvest but before leaf drop.

EVERGREEN TREE CROPS: Such as Avocado, Citrus, Macadamia, Lychee. **Foliar: 3 – 5 L/ha** in a minimum of 600 – 1000L final spray volume. Fertigation, 10 – 20 L/ha. Apply at 14 – 21 day intervals from fruit set to harvest.

FRUITING VEGETABLES: Such as Capsicum, Cucurbits, Eggplant, Tomatoes, Watermelons, Pumpkins. **Foliar: 3 – 8 L/ha** in a minimum of 600 – 1600L final spray volume. **Fertigation: 7 – 15 L/ha.** Apply as required. Wet foliage evenly to drip. When practical use higher (more dilute) water rates. Fertigate during fruiting to replenish nutrients.

LEAFY VEGETABLES: Such as Endive, Fennel Lettuce, Broccoli, Cabbage, Cauliflower, Kale and Herbs. **Foliar: 3 – 6 L/ha** in a minimum of 600 – 1200L final spray volume. **Fertigation: 7 – 10 L/ha.** Apply as required. Every 7 – 14 days from mid crop to harvest. **DO NOT** apply in heat of day.

ROOT VEGETABLES: Such as Beetroot, Carrot, Leek, Onion, Potato, Radish, Sweet Potato. **Foliar: 3 – 8 L/ha** in a minimum of 600 – 1600L final spray volume. **Fertigation: 7 – 10 L/ha.** Apply as required. Wet foliage evenly to drip. When practical use higher (more dilute) water rates.

VINE and BERRY CROPS: Such as Blueberry, Strawberry, Raspberry, Wine and Table Grapes. **Foliar: 2 – 4 L/ha** in a minimum of 400 – 800L final spray volume. **Fertigation: 10 – 15 L/ha.** Apply at 14 day intervals from fruit set onwards. **DO NOT** exceed 2x concentration or 2x hectare rate.

Fertigation rates are dependent on seasonal nutrient demand.

Agitate contents well prior to application.

In hot weather use a more dilute mixture.

The information contained in this Product Information Sheet in respect of the "Product" is indicative only and should not be relied upon as advice or a recommendation. While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information. You use the information at your own risk and should rely on your own independent inquiries and assessments. With the exception of the consumer guarantees provided by the Australian Consumer Law (ACL), all conditions and warranties implied in respect of any information or advice provided by Agrichem about the Product are excluded, and Agrichem does not accept any liability whatsoever (including through misrepresentation or negligence), incurred in connection with your use or reliance upon this Information Sheet. If liability under the ACL cannot be excluded but the Product the subject of the Information Sheet is NOT used for personal, domestic or household use or consumption, Agrichem may (at its election) limit its liability to replacement of the Product, or payment of the cost of acquiring the Product. You must not reproduce this information sheet without written consent from Agrichem©.

NOTE: The suggested rates of application of the Product are designed for typical Australian conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.