

SUPA STAND PHOS®

NPKS 6-10-3-1 + Germination & Root Booster



An effective starter fertiliser for crop establishment, reduce transplant shock, promoting root development and plant growth

BENEFITS OF SUPA STAND PHOS®

- Develops larger, more vigorous root systems through a unique combination of phytoactive compounds and NPK.
- Suitable for various applications to manage the first six – eight weeks of growth.
- Added organic matter to improve soil structure and increase nutrient uptake.
- Improves plant establishment and reduces the incidence of root diseases.

GERMINATION BOOSTER

Supa Stand Phos contains critical phytoactive compounds in a ratio which assists in the stimulation of seed germination and root growth. This ratio will work either as a pre plant dip, furrow injection or foliar spray.

THE IMPORTANCE OF NITROGEN, PHOSPHORUS & POTASSIUM

Nitrogen is the major building block in protein and chlorophyll. It is also essential for lipid and cytoplasm formation. Highly mobile in the plant, it is translocated and utilised in the growing tips.

Phosphorus assists in root development and energy production in plant cells to carry-out vital metabolic functions and nucleic acid biosynthesis. Phosphorus acts as a structural component of nucleic acids and phospholipids which form plant membranes. It is also important in cell division, photosynthesis, sugar and starch formation, energy transfer and movement of carbohydrates. Phosphorus deficiencies are very common in alkaline calcareous and acid soils, due to its binding with calcium in high pH soils and aluminum and iron in acid soils.

Potassium optimises water use efficiency and is the key nutrient to improve crop photosynthesis and sugar production in fruits. Potassium is very important in fruit bearing plants. Potassium regulates the electrolytes and turgidity of plant cells. Potassium occurs in the guard cells of the stomata and is therefore essential in respiration and transpiration. Potassium is required at all growth stages and a lack of potassium cannot be rectified with late applications.

SUPA STAND PHOS[®]

CHARACTERISTICS: pH: 5.5 – 6.5; Specific Gravity: 1.24 – 1.26

AUS Analysis W/W%: 6.2% N, 9.9% P, 3.1% K, 0.98% S, 0.0002% Co, 0.003% Cu, 0.008 Fe, 0.015% Mn, 0.007% Mo, 0.64% Zn
International Analysis W/W%: 5% N, 18.2% P₂O₅, 3% K₂O, 0.78% S, 0.0001% Co, 0.002% Cu, 0.006% Fe, 0.011% Mn, 0.005% Mo, 0.51% Zn

APPLICATION

BROADACRE: Such as Barley, Canola, Cotton, Grain legumes, Maize, Oats, Rice, Sorghum, Triticale, Wheat & Pasture crops. **Foliar: 4 – 7L /ha** in a minimum of 50 - 300 L final spray volume for Ground rigs. **Soil Injection:** Apply via water injection or in-furrow spray at planting or as a foliar from 4 - 6 leaf stage onwards. **Sugar Cane:** Apply at planting- Repeat 2-3 weeks later, if required.

DECIDUOUS TREE CROPS: Such as Apple, Almond, Cherry, Nectarine, Peach, Pear, Pistachio and Walnut. **Foliar: 3 – 5 L/ha** in a minimum of 750 – 1200L final spray volume. Apply 3 sprays, 1st 14 – 21 days post spur burst, 2nd post bloom 3rd 21 days post bloom. **Fertigation: 8 - 15 L/ha. DO NOT apply as a foliar to stonefruits particularly apricots, nectarines and some varieties of peaches during leaf growth.** Can be applied foliar at post harvest but before leaf drop.

EVERGREEN TREE CROPS: Such as Avocado, Banana, Citrus, Macadamia, Mangoes, Lychee. **Foliar: 5 – 10 L/ha** in a minimum of 500 – 1000L final spray volume. Apply to juvenile trees at early establishment - repeat as required. Apply at monthly intervals during active growth period. **Fertigation: 8 - 15 L/ha.**

FRUITING VEGETABLES: Such as Capsicum, Cucurbits, Eggplant, Tomatoes, Watermelons, Pumpkins, Zucchini. **Foliar: 4 – 6 L/ha** in a minimum of 500 – 1000L final spray volume. Apply at emergence or to transplant - repeat at 7 - 10 day intervals as required. **Fertigation: 8 - 15 L/ha. Seedling Dip: 1:100 dilution.**

LEAFY VEGETABLES: Such as Endive, Fennel, Lettuce, Broccoli, Cabbage, Cauliflower, Kale and Herbs. **Foliar: 4 – 6 L/ha** in a minimum of 500 – 1000L final spray volume. **Fertigation: 8 - 15 L/ha** at transplant or emergence. Repeat 7 - 10 days later. **Seedling Dip: 1:100 dilution.**

ROOT VEGETABLES: Such as Beetroot, Carrot, Leek, Onion, Potato, Radish, Sweet Potato. **Foliar: 4 – 6 L/ha** in a minimum of 800 – 1200L final spray volume. **Fertigation: 10 - 15 L/ha.** Apply 1 week after planting - repeat 7 - 10 days later. **Seedling Dip: 1:100 dilution.**

Potato Seed piece dip: 1:3. Apply on top of the seed piece at planting.

VINE and BERRY CROPS: Such as Blueberry, Strawberry, Raspberry, Wine and Table Grapes. **Foliar: 2.5 – 5 L/ha** in a minimum of 1000 – 1600L final spray volume. Apply at 14 day intervals as required. **Fertigation: 10 - 15 L/ha.**

Fertigation rates are dependent on seasonal nutrient demand. Repeat applications may be required.

For nursery plants: drench at 0.1% is highly beneficial.

Agitate contents well prior to application.

WARNING: DO NOT mix or apply with copper based fungicides or apply to crops with copper fungicide residue

The information contained in this Product Information Sheet in respect of the "Product" is indicative only and should not be relied upon as advice or a recommendation. While this Information Sheet has been prepared in good faith, Agrichem does not warrant the accuracy of this information. You use the information at your own risk and should rely on your own independent inquiries and assessments. With the exception of the consumer guarantees provided by the Australian Consumer Law (ACL), all conditions and warranties implied in respect of any information or advice provided by Agrichem about the Product are excluded, and Agrichem does not accept any liability whatsoever (including through misrepresentation or negligence), incurred in connection with your use or reliance upon this Information Sheet. If liability under the ACL cannot be excluded but the Product the subject of the Information Sheet is NOT used for personal, domestic or household use or consumption, Agrichem may (at its election) limit its liability to replacement of the Product, or payment of the cost of acquiring the Product. You must not reproduce this information sheet without written consent from Agrichem©.

NOTE: The suggested rates of application of the Product are designed for typical Australian conditions and should be used as a guide only. Each farmer's climatic conditions, water quality, soil types, application processes and practices may differ and therefore necessitate corrections to ensure optimum results. Good agricultural practice requires that application be avoided under extreme weather conditions such as temperatures over 28°C, high humidity, frost, rain etc. It is recommended that when applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total spray. Where possible, it is recommended that regular leaf tests are conducted to determine actual plant nutrient availability during each growth cycle. Soil tests at least once per year are essential.